

## Aryabhata's trig table Math 105 History of Mathematics Prof. D. Joyce, Spring 2015

Aryabhata (b. 476) included a table of sines in his Aryabhatiya and a rule for constructing that table of sines. For Aryabhata, a sine was a half-chord in a circle of radius 3438 (the same radius Hipparchus had used centuries earlier). Thus, Aryabhata's sine for an angle  $\theta$  equals 3438 sin  $\theta$ . His table is given in increments of 3° 45′ for angles strictly between 0° and 90°, but only increases in sines are given.

Stanza I, 10. The twenty-four sine [differences] reckoned in minutes of arc are 225, 224, 222, 219, 215, 210, 205, 199, 191, 183, 174, 164, 154, 143, 131, 119, 106, 93, 79, 65, 51, 37, 22, 7.

If we denote these differences by  $d_1, d_2, \ldots, d_{24}$ , and their sums by  $s_1 = d_1, s_2 = d_1 + d_2, \ldots, s_{24} = d_1 + d_2 + \cdots + d_{24}$ , then the sums are the sines of various angles. The first entry, 225, gives  $s_1 = 3438 \sin 3^{\circ} 45'$ . Add to that the second,  $d_2 = 224$ , to give  $s_2 = 449 = 3438 \sin 7^{\circ} 30'$ . Add to that the third,  $d_3 = 222$ , to give  $s_3 = 671 = 3438 \sin 11^{\circ} 15'$ . And so forth.

Thus, if you've memorized the stanza, you can construct a table of sines for trigonometry since you can easily compute a sine from the previous sine and the sine difference:  $s_n = s_{n-1} + d_n$ .

In a later stanza, Aryabhata gives a rule for constructing the twenty-four sine differences. This stansa tells how to compute the differences  $d_n$ .

Stanza II, 12. By what number the last sine [difference] is less than the first sine, and by the quotient obtained by dividing the sum of the preceding sine [differences] by the first sine, by the sum of these two

quantities the following sine [differences] are less than the first sine.

As an equation, this rule says

$$(d_1 - d_{n-1}) + \frac{d_1 + d_2 + \dots + d_{n_1}}{d_1} = d_1 - d_n$$

or, more simply,

$$d_n = d_n - s_{n-1}/225.$$

Below is an table of the values. It only depends on the two equations,  $s_n = s_{n-1} + d_n$  and  $d_n = d_n - s_{n-1}/225$ , and the values in the first line. The numbers in the last column,  $d_n$ , are usually rounded down to the nearest integer, but sometimes rounded up to the next integer.

$\underline{}$	$d_n$	$s_n$	$s_n/225$
1	225	225	225/225 = 1.0
2	224	449	449/225 = 2.0
3	222	671	671/225 = 3.0
4	219	890	890/225 = 4.0
5	215	1105	1105/225 = 4.9
6	210	1315	1315/225 = 5.8
7	205	1520	1520/225 = 6.8
8	199	1719	1719/225 = 7.6
9	191	1910	1910/225 = 8.5
10	183	2093	2093/225 = 9.3
11	174	2267	2267/225 = 10.1
12	164	2431	2431/225 = 10.8
13	154	2585	2585/225 = 11.5
14	143	2728	2728/225 = 12.1
15	131	2859	2859/225 = 12.7
16	119	2978	2978/225 = 13.2
17	106	3084	3084/225 = 13.7
18	93	3177	3177/225 = 14.1
19	79	3256	3256/225 = 14.5
20	65	3321	3321/225 = 14.8
21	51	3372	3372/225 = 14.9
22	37	3409	3409/225 = 15.2
23	22	3431	3431/225 = 15.2
24	7	3438	•

Math 105 Home Page at

http://math.clarku.edu/~djoyce/ma105/